

RR Lyrae Stars in the Far Ultraviolet: GALEX Observations Compared with Theoretical Predictions

S.E. Browne, J.M. Wheatley, B.Y. Welsh (University of California Berkeley), M. Seibert, K. Heafield (Caltech), R.M. Rich (UCLA) and the GALEX Science Team

The GALEX Mission

- GALEX is a NASA Small Explorer satellite, launched in April 2003, and is currently surveying the sky at UV wavelengths from 1350-2800Å.
- Several imaging surveys are in progress:
 - AIS: All Sky Survey, 40000 degrees² to AB=20.5 mag
 - MIS: Medium Imaging Survey, 1,000 degrees² to AB=23 mag
 - DIS: Deep Imaging Survey, 80 degrees² to AB=25 mag
 - NGS: Nearby Galaxy Survey, 240 degrees² to AB=23 mag
- Photon-counting detectors enable high time-resolution (<0.01s) UV photometric observations.
- Independent detectors simultaneously observe at FUV (1350-1750Å) and NUV (1750-2800Å)
- Public releases of GALEX data are available at galex.stsci.edu, at the Multi-Mission Archive at the Space Telescope Science Institute (MAST).

Standard Candles in the Ultraviolet

- RR Lyrae stars in crowded fields in the cores of globular clusters and in the Magellanic Clouds may be easier to detect in the UV than in the visible
- RR Lyraes are far less luminous in the FUV than in the visible, making them hard to detect in distant galaxies

GALEX RR Lyrae Observations

- The wide field of view (1.2 degrees), high sensitivity (AB~23 mag in 1500 s), and low background enable GALEX to serendipitously detect many variable and transient sources during its surveys
- In the MIS and DIS surveys, GALEX makes repeat visits to star fields: this has proven ideal for producing high-quality RR-Lyrae light curves at FUV and NUV wavelengths
- We illustrate GALEX light curves for six RR Lyrae ab-type variables that have many observations well distributed in phase (central panels).
- RR Lyraes vary by 4-6 magnitudes in FUV, and by 2-3 mag in NUV, far more than the ~1 mag variations at visible wavelengths

The Pulsation Cycle

- All ab-type RR Lyraes have a minimum atmospheric temperature of ~6000K, occurring between phases .5 and .85
- At maximum light, the peak temperature can reach anywhere from 7000K to 7800K, depending on the mass of the star
- The star's radius reaches a minimum at phase 0.85, just before the rapid brightness rise to maximum, which accounts for the slight dip at this phase
- Surface gravity varies as a function of the star's radius, and also due to radial acceleration during expansion and contraction in the pulsation cycle

Kurucz Model Atmospheres in the FUV

- Kurucz (2003) has computed ultraviolet stellar spectra over a wide range of metallicities, temperatures, and surface gravities, using the improved opacities of Iglesias, et al. (1996).
- In the GALEX FUV band (1350-1800Å), Kurucz predicted fluxes are highly sensitive to metallicity and temperature in the 6000-7500K range, but far less sensitive to surface gravity (log g, cgs units).
- This contrasts with longer wavelengths (4000-10000Å), where the predicted RR Lyrae flux is largely insensitive to metallicity.
- We explore a range of temperature and metallicity to fit the observed GALEX and visible light curves




Figure 1. RR Lyrae light curves in NUV (1800-2800Å, open circles) and FUV (1350-1800Å, filled circles), with one sigma error bars. Amplitudes are ~2.5 mag in NUV and 5-6 mag in FUV. Overlaid are Kurucz model atmosphere light curves (see below) for [Fe/H]=0.0 (short dash), [Fe/H]=-1.0 (long dash) and [Fe/H]=-1.75 (solid line).


Fitting Models to the GALEX Light Curves

- We begin with a temperature variation of 5900-7300K, deduced from visible photometry of ROTSE-I J143753.84+345924.8 (lower right in Fig. 1) by Wheatley et al. (2005).
- Metallicities ranging from [Fe/H]=0.0 to -1.75 are explored to find the best fit to the ultraviolet light curves
- Lower metallicities produce dramatically larger amplitudes in the FUV light curves
- The observed 5-6 magnitude FUV variations are consistent with metallicities of -1.0 to -1.5
- Model light curves are poorly constrained by UV data alone
- Our continuing Lick Observatory photometry program will pinpoint the temperature variations of these stars, allowing more accurate UV-derived metallicities

Reference: Wheatley, J.M. et al. 2005, Large-Amplitude Ultraviolet Variations in the RR Lyrae Star ROTSE-I J143753.84+345924.8. *Astrophysical Journal*, Volume 619, Issue 1, pp. L123-L126.


GALEX Aperture Curves of Growth

Several stars with a small range of NUV or FUV magnitudes in GALEX fields are examined to produce aperture growth curves. Curves with a small variance are compared to curves for other stars within a selected magnitude range in different exposures to produce a time series to check for variability.




Visible Observations of GALEX Variables

If only a few GALEX observations exist, as here, visible observations (at the Lick Observatory) are made to confirm & determine the nature and parameters of the variability.



From Visible Observations, A Light Curve Emerges

In this case, a light curve emerged through a "least string" analysis of the visible Lick data. The light curve suggests this source is an RR Lyrae type "c" or a high-amplitude Delta Scuti star.



Conclusions

- GALEX FUV light curves with amplitudes of 5-6 magnitudes are consistent with Kurucz model atmosphere predictions.
- Ongoing optical V-band photometry at the Lick Observatory will better constrain the metallicities derived from GALEX data
- New UV variable star candidates can be discovered in GALEX observations by a comparative curve-of-growth photometric analysis
- Future GALEX observations may produce FUV and NUV light curves of other kinds of pulsating variables, such as δ Scuti stars